

Inclusive and "exclusive" cross sections in the regime of multiple parton collisions

G. Calucci* and D. Treleani†

*Dipartimento di Fisica Teorica dell'Università di Trieste and INFN, Sezione di Trieste,
Strada Costiera 11, Miramare-Grignano, I-34014 Trieste, Italy.*

The large luminosity and the increased kinematical domain accessible at the LHC will allow to isolate large numbers of events with multiple parton collisions. Interestingly, the hadron is probed in different points simultaneously in the process, which allows to obtain information on the correlations between partons in the hadron structure. The whole experimentally accessible information on multiple parton collisions consists in the probability distribution of the elementary interactions, while the inclusive cross sections, usually considered in large p_t processes, acquire a precise statistical meaning as moments of the multiplicity distribution. Conversely different and more "exclusive" cross sections become experimentally viable, providing complementary information on the interaction dynamics. The matter is discussed in details in the case of hadron-hadron collisions and the role of parton correlations is outlined both in the inclusive and in the "exclusive" cross sections.

PACS numbers: 11.80.La; 12.38.Bx; 13.85.Hd; 13.87.-a

Keywords: Multiple scattering, Perturbative calculations, Inelastic scattering. Multiple production of jets

1. INTRODUCTION

The overall features of the minimum bias and of the underlying event in hard hadronic interactions cannot be reproduced without implementing multiple parton interactions (MPI) in the event generator[1][2][3][4]. On the other hand, the direct measurement of MPI requires identifying the final fragments of the multiple process, to reconstruct the kinematics at the parton level. The sizable reduction of statistics, caused by the request of large momentum exchange in each hard interaction, has restricted considerably up to now the possibilities of a direct study of the phenomenon and measurements have been possible only in very limited phase space intervals[5][6][7][8]. The large energy in the pp c.m. system may however justify the expectation that, at the Large Hadron Collider (LHC), MPI will not be important only to describe the properties of the minimum bias and of the underlying event, giving a significant contribution also, as an example, to the cross section to produce jets with large p_t .

A preliminary requirement for the study of MPI in large p_t jet production is the separation of the contribution of multiple radiation, as the two mechanisms can produce the same final state. When looking at the properties of the minimum bias and of the underlying event the separation between the two sources is rather ambiguous. In the case of direct observation of the large p_t scattered partons, one may however expect that the separation of the contribution of MPI form

*Electronic address: giorgio.calucci@ts.infn.it

†Electronic address: daniele.treleani@ts.infn.it

the leading single scattering perturbative Quantum Chromo Dynamics (QCD) mechanism might be done more confidently. On one side hard radiation induces an azimuthal correlation between the produced jets, which is absent in the case of jets produced through MPI, on the other the two mechanisms have a very different dependence on the lower value of the transverse momentum introduced to observe the jets and on the c.m. energy, being the MPI a power correction to the leading QCD single scattering mechanism. Azimuthal correlations were in fact successfully used by CDF to isolate and measure double parton collisions at Fermilab and the statement made by the experiment, that some 17% of their sample of double scattering events was contaminated by triple scatterings[8], justifies the hope that the larger energy available at the LHC will allow to isolate events with triple and perhaps also with quadruple parton interactions without much ambiguity.

One should nevertheless point out that, while in the case of double collisions the separation of hard radiation and of the MPI contribution may be relatively simple, the best strategy to isolate the contribution of multiple collisions, when the number of scatterings is large, may not be readily identified. Although it might still be relatively simple, at least for triple scatterings, in the case of particular channels, like W production plus jets or Drell-Yan plus jets, with the jets in particular kinematical configurations. In more general cases, the strategy to separate the contribution of MPI from the contribution of the leading single QCD production mechanism deserves a dedicated study and will not be addressed in the present paper. Rather we will assume that the increased parton luminosity and the sizably higher value of the momentum exchanged in a typical partonic interaction at the LHC will allow the direct identification of a sizable number of events, not only with double but also with triple and perhaps quadruple parton interactions.

Interestingly, one may expect that a detailed study of the phenomenon will allow to access several features of the hadron structure still unexplored and most likely unexpected. The anomalously small value of the scale factor, characterizing the measurement of double parton interactions by CDF at Tevatron[7][8][9], and the need to use a parton density much different as compared with the hadron form factor, in the actual Montecarlo generators[1][3], are indications that unpredicted features of the hadronic structure may play a non secondary role in the phenomenon.

A general frame to identify and organize the novel information in a systematic way is hence highly desirable. The present paper represents an attempt to provide such a frame. One should stress that our goal is by no means to provide a scheme for a general description of the overall inelastic event, so our aim is very different in comparison with Montecarlo models. Our final states consists only of the large p_t partons generated by MPI, where we assume that one will be able to recognize and count the number of interactions in some phase space windows. In addition, we are interested only in a small fraction of all inelastic events and only in few particular final state observables (like for example the jet's momenta). Our aim is to show how such quantities are linked to the correlations of the partons in the hadron structure, while remaining as general as possible. So we will not attempt to compare various ideas about correlations, which have been explicitly implemented, in particular in Montecarlo models. Rather we will try to identify observables which have a direct link with correlations and to establish sum rules which, allowing to control the consistency of the scheme at a given order in the number of MPI, will help to disentangle the effects of correlations from the effects of unitarization.

A preliminary comment concerns the hard cross sections accessible in the regime of MPI. As

it will be discussed in the next sections, in the case of hadron-hadron interactions the integrated inclusive cross sections of the QCD-parton model are essentially the moments of the distribution in the number of collisions. The whole information of a probabilistic distribution is either given by the set of the different terms of the distribution or, alternatively, by the set of its moments. One may hence consider two different sets of cross sections, the inclusive cross sections, given by the moments of the distribution, and the "exclusive" cross sections, given by the probabilities of the different terms of the distribution in the number of collisions.

The two different sets of cross sections correspond to two different criteria of selection of events. In the case of the inclusive cross section of a K -parton scattering, one needs to collect all events where there are *at least* K -parton collisions in a given final state phase space interval. All events with N -parton collisions, where $N \geq K$, hence contribute with a multiplicity factor $\binom{N}{K}$. On the other hand the "exclusive" cross section of a K -parton scattering process is measured by selecting all events with *only* K -parton collisions in the phase space interval of interest. The cross sections called here "exclusive" are in fact partially inclusive cross sections, since one sums over all partons outside the given phase space interval and on the soft fragments.

By measuring the inclusive cross sections, which are related to the moments of the distribution, one has the advantage of approaching the problem in a way ordered in complexity. In the case of the "exclusive" cross sections, each term is on the contrary related to the whole set of the multiple scattering series, since each term contains the probability of no interaction, which carries information on the whole series of multiple collisions. In other words each "exclusive" cross section requires, at least in principle, the knowledge of an infinite set of non perturbative inputs to be evaluated. On the other hand the "exclusive" cross sections are well defined experimental quantities, characterizing the regime of hard interactions. It is hence worth trying to discuss also the "exclusive" cross sections in the framework of perturbation theory.

The simplest case of "exclusive" cross section is the hard cross section itself, defined as the contributions to the inelastic cross section of all events with at least one hard partonic interaction. In the case of hadron-hadron collisions all terms with correlations can be re-summed, when only two-body parton correlations are taken into account[9]. As a result of resummation the effects of the correlation terms is rather indirect in the hard cross section, in such a way that the information on parton correlations cannot be disentangled in a simple way. Rather obviously a similar property holds for all "exclusive" cross sections.

Isolating and organizing the correlation terms in order of growing importance is nevertheless possible also in the case of the "exclusive" cross sections. As a consequence of incoherence between the different partonic interactions, MPI are in fact described by a (different) probabilistic distribution in each phase space interval and one may reduce or increase the average number of partonic collisions by restricting or enlarging the interval in transverse momenta and in rapidity where final states are observed. It is hence always possible to give an operational meaning to an expansion in the number of collisions. The expansion of the "exclusive" cross sections in number of elementary collisions allows to isolate the different correlation terms and to obtain interesting consistency relations between inclusive and "exclusive" cross sections.

The paper is organized as follows. After recalling the main features of the simplest Poissonian model, which represents the basis of most of the practical implementations of multiple parton collisions, we will generalize the approach by including parton correlations in the picture of the interaction through a functional approach. Keeping only two-body parton correlations into

account, explicit expressions will hence be obtained both for inclusive and "exclusive" cross sections, in the latter case by expanding in the number of parton collisions. A simple and experimentally testable connection between the two quantities, at the lowest non trivial order in the number of collisions, will be finally derived.

2. THE SIMPLEST POISSONIAN MODEL

A standard way to introduce MPI in Montecarlo codes is to assume a Poissonian distribution of multiple parton collisions, with average number depending on the value of the impact parameter. The approach was introduced long ago[10][11] and critically re-discussed recently[12]. Of course the Poissonian is only a starting point for the Montecarlo generators, which are in fact much more elaborated. Both the JIMMY[13] and PYTHIA[1] include momentum correlations and the newest variant of PYTHIA[14] imposes also flavor conservation. In the present section we discuss the simplest option, namely the case where no correlation is taken into account in a hadronic interaction at a given value of the impact parameter. Correlations will be introduced in a rather general way in the following section.

One may start by introducing the three dimensional parton density $D(x, b)$, namely the average number of partons with a given momentum fraction x and with transverse coordinate b (the dependence on flavor and on the resolution of the process is understood) and one makes the simplifying assumption $D(x, b) = G(x)f(b)$, with $G(x)$ the usual parton distribution function and $f(b)$ normalized to one. The inclusive cross section for large p_t parton production σ_S may hence be expressed as

$$\sigma_S = \int_{p_t^c} G(x)\hat{\sigma}(x, x')G(x')dx dx' = \int_{p_t^c} G(x)f(b)\hat{\sigma}(x, x')G(x')f(b-\beta)d^2b d^2\beta dx dx' \quad (1)$$

where p_t^c is a cutoff introduced to distinguish hard and soft parton collisions and β the hadronic impact parameter. The expression allows a simple geometrical interpretation. Given the large momentum exchange, which localizes the partonic interaction inside the overlap volume of the two hadrons, one may identify with b and $b - \beta$ the transverse coordinates of the two colliding partons while β is the impact parameter of the hadronic collision.

Neglecting all correlations in the multi-parton distributions, the inclusive cross section for a double parton scattering σ_D is analogously expressed by

$$\begin{aligned} \sigma_D &= \frac{1}{2!} \int_{p_t^c} G(x_1)f(b_1)\hat{\sigma}(x_1, x'_1)G(x'_1)f(b_1-\beta)d^2b_1 dx_1 dx'_1 \times \\ &\quad \times G(x_2)f(b_2)\hat{\sigma}(x_2, x'_2)G(x'_2)f(b_2-\beta)d^2b_2 dx_2 dx'_2 d^2\beta \\ &= \int \frac{1}{2!} \left(\int_{p_t^c} G(x)f(b)\hat{\sigma}(x, x')G(x')f(b-\beta)d^2b dx dx' \right)^2 d^2\beta \end{aligned} \quad (2)$$

which allows an analogous geometrical interpretation. The expression may be readily generalized to the case of the inclusive cross section for N -parton scatterings σ_N :

$$\sigma_N = \int \frac{1}{N!} \left(\int_{p_t^c} G(x)f(b)\hat{\sigma}(x, x')G(x')f(b-\beta)d^2b dx dx' \right)^N d^2\beta \quad (3)$$

The cross sections are divergent for $p_t^c \rightarrow 0$. The unitarity problem is solved by normalizing the integrand which, being dimensionless, may be understood as the probability to have a N th parton collision process in a inelastic event. One may hence introduce $P_N(\beta)$, the probability of having N parton collisions in a hadronic interaction at impact parameter β :

$$P_N(\beta) \equiv \frac{(\sigma_S F(\beta))^N}{N!} e^{-\sigma_S F(\beta)}, \quad \int_{p_t^c} G(x) f(b) \hat{\sigma}(x, x') G(x') f(b - \beta) d^2 b dx dx' \equiv \sigma_S F(\beta) \quad (4)$$

By summing all probabilities one obtains the hard cross section σ_{hard} , namely the contribution to the inelastic cross section due to all events with *at least* one parton collision with momentum transfer greater than the cutoff p_t^c :

$$\sigma_{hard} = \sum_{N=1}^{\infty} \int P_N(\beta) d^2 \beta = \sum_{N=1}^{\infty} \int d^2 \beta \frac{(\sigma_S F(\beta))^N}{N!} e^{-\sigma_S F(\beta)} = \int d^2 \beta [1 - e^{-\sigma_S F(\beta)}] \quad (5)$$

Notice that σ_{hard} is finite in the infrared limit, which allows to express the inelastic cross section as $\sigma_{inel} = \sigma_{soft} + \sigma_{hard}$ with σ_{soft} the soft contribution, the two terms σ_{soft} and σ_{hard} being defined through the cutoff in the momentum exchanged at parton level, p_t^c .

As mentioned the inclusive cross sections are related to the moments of the distribution. The property is immediate in the case of the Poissonian model. Working out the average number of collisions one in fact obtains:

$$\langle N \rangle \sigma_{hard} = \int d^2 \beta \sum_{N=1}^{\infty} N P_N(\beta) = \int d^2 \beta \sum_{N=1}^{\infty} \frac{N [\sigma_S F(\beta)]^N}{N!} e^{-\sigma_S F(\beta)} = \int d^2 \beta \sigma_S F(\beta) = \sigma_S \quad (6)$$

which is the single scattering inclusive cross section, while, more in general, one may write:

$$\begin{aligned} \frac{\langle N(N-1) \dots (N-K+1) \rangle}{K!} \sigma_{hard} &= \int d^2 \beta \sum_{N=1}^{\infty} \frac{N(N-1) \dots (N-K+1)}{K!} P_N(\beta) \\ &= \int d^2 \beta \frac{1}{K!} [\sigma_S F(\beta)]^K = \sigma_K \end{aligned} \quad (7)$$

A few features of the model are worth to be pointed out.

- The cross sections of the two sets, inclusive and "exclusive" (the latter being the contributions to σ_{hard} due to the different numbers of collisions) are expressed fully explicitly in terms of the same quantity, the average number of collisions at a fixed impact parameter $\sigma_S F(\beta)$. In a more general case, when correlations are taken into account, the two sets of cross sections can provide a complementary information on the multi parton structure of the hadron.
- All inclusive cross sections are divergent in the infrared limit. The relation with the multiplicity of collisions shows that the divergence is due to the moments. In other words the cause of the divergence is the number of collisions, which become very large at low p_t .
- While all inclusive cross sections become increasingly large at low p_t , due to the increasingly large number of parton interactions, all "exclusive" cross section where the number of participating hard partons is kept fixed become, on the contrary, smaller and smaller for the same reason.

- In the model one assumes that the transverse hadron size is the same in each interaction, so one is identifying the hadron with its average configuration. Hadronic diffraction shows however that hadron fluctuations are not a negligible effect also at LHC energies[15][16]. It was moreover pointed out that, most likely, the importance of hadronic fluctuations grows when the number of partonic collisions is large[17][18].

3. A FUNCTIONAL APPROACH

MPI can be discussed in very general terms by a functional formalism[19][9].

Let $W_n(u_1 \dots u_n)$ be the exclusive n -parton distributions, namely the probabilities to have the hadron in a configuration with n partons with coordinates u_i , which represent the variables (b_i, x_i) , being b the transverse partonic coordinate and x the corresponding fractional momentum. The scale for the distributions is given by the cut off p_t^{min} that defines the separation between soft and hard collisions. The distributions are symmetric in the variables u_i . One may introduce the generating functional:

$$\mathcal{Z}[J] = \sum_n \frac{1}{n!} \int J(u_1) \dots J(u_n) W_n(u_1 \dots u_n) du_1 \dots du_n. \quad (8)$$

The conservation of the probability implies the normalization condition $\mathcal{Z}[1] = 1$.

The probabilities of the various configurations, namely the exclusive distributions, are the coefficients of the expansion of $\mathcal{Z}[J]$ for $J = 0$. The coefficients of the expansion of $\mathcal{Z}[J]$ for $J = 1$ give the many body densities, i.e. the inclusive distributions:

$$D_1(u) = \left. \frac{\delta \mathcal{Z}}{\delta J(u)} \right|_{J=1}, \quad D_2(u_1, u_2) = \left. \frac{\delta^2 \mathcal{Z}}{\delta J(u_1) \delta J(u_2)} \right|_{J=1} \dots \quad (9)$$

Correlations, which describe how much the distribution deviates from a Poissonian, are obtained by the expansion of the logarithm of the generating functional, $\mathcal{F}[J] \equiv \ln \mathcal{Z}[J]$, for $J = 1$:

$$\mathcal{F}[J] = \int D_1(u) [J(u) - 1] du + \sum_{n=2}^{\infty} \frac{1}{n!} \int C_n(u_1 \dots u_n) [J(u_1) - 1] \dots \dots [J(u_n) - 1] du_1 \dots du_n \quad (10)$$

Obviously one has $\mathcal{F}[1] = 0$ and, in the Poissonian case, $C_n \equiv 0, n \geq 2$.

Given the general expressions for the multiparton distributions one can write down an expression for the semi-hard cross section which takes into account of all possible MPI:

$$\begin{aligned} \sigma_{hard} = \int d\beta \int & \sum_n \frac{1}{n!} \frac{\delta}{\delta J(u_1)} \dots \frac{\delta}{\delta J(u_n)} \mathcal{Z}_A[J] \\ & \times \sum_m \frac{1}{m!} \frac{\delta}{\delta J'(u'_1 - \beta)} \dots \frac{\delta}{\delta J'(u'_m - \beta)} \mathcal{Z}_B[J'] \end{aligned}$$

$$\times \left\{ 1 - \prod_{i=1}^n \prod_{j=1}^m [1 - \hat{\sigma}_{i,j}(u, u')] \right\} \prod dud u' \Big|_{J=J'=0} \quad (11)$$

Here β is the impact parameter between the two interacting hadrons A and B and $\hat{\sigma}_{i,j}$ is the probability for the parton i (of A) to have an hard interaction with the parton j (of B).

The cross section is obtained by summing all contributions due to all different hadronic configurations (the sums over n and m). For each pair of values n and m , one has a contribution to σ_{hard} when at least one hard interaction takes place (whose probability is represented by the term in curly brackets in the equation above).

The cross section, which is analogous to the expression of the inelastic nucleus-nucleus cross section in the Glauber model[20], takes into account both disconnected interactions (which imply $n = m$) and rescatterings (when $n \neq m$).

A much simpler expression may be obtained by neglecting all rescatterings. To that purpose one may write the term in curly brackets as:

$$S \equiv 1 - \exp \sum_{ij} \ln(1 - \hat{\sigma}_{ij}) = 1 - \exp \left[- \sum_{ij} \left(\hat{\sigma}_{ij} + \frac{1}{2} \hat{\sigma}_{ij} \hat{\sigma}_{ij} + \dots \right) \right] \quad (12)$$

rescatterings correspond to repeated indices. Only the first term of the expansion of the logarithm hence contributes and all rescatterings are removed by the substitution:

$$S \Rightarrow 1 - \exp \sum_{ij} (\hat{\sigma}_{ij}) \Rightarrow \sum_{ij} \hat{\sigma}_{ij} - \frac{1}{2} \sum_{ij} \sum_{k \neq i, l \neq j} \hat{\sigma}_{ij} \hat{\sigma}_{kl} \dots \quad (13)$$

The resulting cross section is expressed in a compact way

$$\begin{aligned} \sigma_{hard}(\beta) &= \exp(\partial) \cdot \exp(\partial') \left[1 - \exp(-\partial \cdot \hat{\sigma} \cdot \partial') \right] \mathcal{Z}_A[J] \mathcal{Z}_B[J'] \Big|_{J=J'=0} \\ &= \left[1 - \exp(-\partial \cdot \hat{\sigma} \cdot \partial') \right] \mathcal{Z}_A[J] \mathcal{Z}_B[J'] \Big|_{J=J'=1} \end{aligned} \quad (14)$$

where all convolutions are understood. In the simplest non-trivial case all correlations C_n with $n > 2$ can be neglected and the cross section is more explicitly expressed by:

$$\begin{aligned} \sigma_{hard}(\beta) &= \left[1 - \exp \left\{ - \int dud u' \partial_J \hat{\sigma}(u, u') \partial_{J'} \right\} \right] \\ &\cdot \exp \left\{ \int D_A(u) J(u) du + \frac{1}{2} \int C_A(u, v) J(u) J(v) dudv \right\} \\ &\cdot \exp \left\{ \int D_B(u) J(u) du + \frac{1}{2} \int C_B(u, v) J(u) J(v) dudv \right\} \Big|_{J=J'=0} \end{aligned} \quad (15)$$

which can be worked out explicitly. One in fact obtains[19][9]

$$\sigma_{hard}(\beta) = 1 - \exp \left[- \frac{1}{2} \sum_n a_n - \frac{1}{2} \sum_n b_n / n \right] \quad (16)$$

where

$$a_n = (-1)^{n+1} \int D_A(u_1) \hat{\sigma}(u_1, u'_1) C_B(u'_1, u'_2) \hat{\sigma}(u'_2, u_2) C_A(u_2, u_3) \dots \dots \hat{\sigma}(u_n, u'_n) D_B(u'_n) \prod_{i=1}^n du_i du'_i \quad (17)$$

and

$$b_n = (-1)^{n+1} \int C_A(u_n, u_1) \hat{\sigma}(u_1, u'_1) C_B(u'_1, u'_2) \dots \dots C_B(u'_{n-1}, u'_n) \hat{\sigma}(u'_n, u_n) \prod_{i=1}^n du_i du'_i \quad (18)$$

Notice that if one works out the moments of the distribution in the number of collisions one obtains the same result of the simplest Poissonian model. From Eq.14 one may express the hard cross section as a sum of MPI:

$$\begin{aligned} \sigma_{hard}(\beta) &= \left[1 - \exp(-\partial \cdot \hat{\sigma} \cdot \partial') \right] \mathcal{Z}_A[J] \mathcal{Z}_B[J'] \Big|_{J=J'=1} \\ &= \sum_{N=1}^{\infty} \frac{(\partial \cdot \hat{\sigma} \cdot \partial')^N}{N!} e^{-\partial \cdot \hat{\sigma} \cdot \partial'} \mathcal{Z}_A[J] \mathcal{Z}_B[J'] \Big|_{J=J'=1} \end{aligned} \quad (19)$$

The average number of collisions hence is

$$\begin{aligned} \langle N \rangle \sigma_{hard}(\beta) &= \sum_{N=1}^{\infty} \frac{N (\partial \cdot \hat{\sigma} \cdot \partial')^N}{N!} e^{-\partial \cdot \hat{\sigma} \cdot \partial'} \mathcal{Z}_A[J] \mathcal{Z}_B[J'] \Big|_{J=J'=1} \\ &= \partial_{J_1} \cdot \hat{\sigma} \cdot \partial_{J'_1} \sum_{N=0}^{\infty} \frac{(\partial \cdot \hat{\sigma} \cdot \partial')^N}{N!} e^{-\partial \cdot \hat{\sigma} \cdot \partial'} \mathcal{Z}_A[J] \mathcal{Z}_B[J'] \Big|_{J=J'=1} \\ &= (\partial_{J_1} \cdot \hat{\sigma} \cdot \partial_{J'_1}) \mathcal{Z}_A[J] \mathcal{Z}_B[J'] \Big|_{J=J'=1} \\ &= \int D_A(x_1; b_1) \hat{\sigma}(x_1 x'_1) D_B(x'_1; b_1 - \beta) dx_1 dx'_1 d^2 b_1 \equiv \sigma_S(\beta) \end{aligned} \quad (20)$$

where now $\hat{\sigma}(x_1 x'_1)$ is the parton-parton cross section integrated with $p_t > p_t^c$ since, given the localization of the interactions in transverse space, the parton-parton interaction probability has been treated as a δ as a function of the transverse coordinates: $\hat{\sigma}(u, u') = \hat{\sigma}(x, x') \delta(\mathbf{b} - \mathbf{b}')$. Analogously one obtains

$$\begin{aligned} \frac{\langle N(N-1) \rangle}{2!} \sigma_{hard}(\beta) &= \frac{1}{2!} \int D_A(x_1 x_2; b_1 b_2) \hat{\sigma}(x_1 x'_1) \hat{\sigma}(x_2 x'_2) \\ &\quad \times D_B(x'_1 x'_2; b_1 - \beta, b_2 - \beta) dx_1 dx'_1 d^2 b_1 dx_2 dx'_2 d^2 b_2 \\ &\equiv \sigma_D(\beta) \end{aligned} \quad (21)$$

and in general

$$\begin{aligned}
& \frac{\langle N(N-1)\dots(N-K+1) \rangle}{K!} \sigma_{hard}(\beta) \\
&= \frac{1}{K!} \int D_A(x_1 \dots x_K; b_1 \dots b_K) \hat{\sigma}(x_1 x'_1) \dots \hat{\sigma}(x_K x'_K) \\
&\quad \times D_B(x'_1 \dots x'_K; b_1 - \beta \dots b_K - \beta) dx_1 dx'_1 d^2 b_1 \dots dx_K dx'_K d^2 b_K \\
&\equiv \sigma_K(\beta)
\end{aligned} \tag{22}$$

which hence shows that *when rescatterings are neglected*, for any choice of multiparton distributions, the inclusive cross sections are given by the moments of the distribution in the number of collisions.

4. "EXCLUSIVE" CROSS SECTIONS

As discussed in the previous section, the inclusive cross sections are basically the moments of the distribution of the number of collisions. The moments contain the whole information on the distribution in the number of collisions ordered in complexity: The average number is the most basic information on the distribution and is given by the single scattering inclusive cross section of the QCD parton model which, precisely for this reason, represents also the simplest theoretical quantity to evaluate. Its expression is in fact particularly simple, the convolution of the parton distributions (which represent the average number of partons in the hadron in a given kinematical configuration) and the single scattering partonic cross section.

Analogously the K -parton scattering inclusive cross section gives the K th moment of the distribution in the number of collisions and is related directly to the K -body parton distribution of the hadron structure.

As already pointed out, a way alternative to the set of moments, to provide the information of the distribution, is represented by the set of the different terms of the probability distribution of multiple collisions. Correspondingly, in addition to the set of the inclusive cross sections, one may consider the set of the "exclusive" cross sections. As discussed in the previous section the case where only two-body parton correlations are taken into account can be worked out in details. In the following we will hence work out explicit expression for the single and for the double parton scattering "exclusive" cross sections.

To obtain the distribution of multiple interactions one may start from the partonic interaction probability

$$1 - \prod_{i,j=1}^n (1 - \hat{\sigma}_{ij}) \tag{23}$$

where in the product each index assumes a given value only once, in such a way that possible re-interactions are not included. The probability of a single interaction may hence be expressed as

$$\left(-\frac{\partial}{\partial g} \right) \prod_{i,j=1}^n (1 - g\hat{\sigma}_{ij}) \Big|_{g=1} = \sum_{kl} \hat{\sigma}_{kl} \prod_{ij \neq kl}^n (1 - g\hat{\sigma}_{ij}) \Big|_{g=1} \tag{24}$$

while the probability for a double collisions is

$$\frac{1}{2!} \left(-\frac{\partial}{\partial g} \right)^2 \prod_{i,j=1}^n (1 - g\hat{\sigma}_{ij}) \Big|_{g=1} = \frac{1}{2!} \sum_{kl} \sum_{rs} \hat{\sigma}_{kl} \hat{\sigma}_{rs} \prod_{ij \neq kl, rs}^n (1 - g\hat{\sigma}_{ij}) \Big|_{g=1} \quad (25)$$

The expressions for the single and double "exclusive" integrated cross sections are

$$\left(-\frac{\partial}{\partial g} \right) e^{-X(g)} \Big|_{g=1} = X'(g) e^{-X(g)} \Big|_{g=1} \quad (26)$$

$$\frac{1}{2!} \left(-\frac{\partial}{\partial g} \right)^2 e^{-X(g)} \Big|_{g=1} = \frac{1}{2!} \left\{ [X'(g)]^2 - X''(g) \right\} e^{-X(g)} \Big|_{g=1} \quad (27)$$

where $X = \frac{1}{2}(\sum a_n + \sum b_n/n)$ with a_n and b_n given by Eq.'s 17 and 18.

The physical description of the process is purely probabilistic. As a consequence, the hard cross section may be unitarized in each given phase space volume, which hence defines also all integration limits.

By restricting the rapidity window or the p_t interval one may control the importance of the terms with different powers of $\hat{\sigma}$. In a small window only the terms linear in $\hat{\sigma}$ are relevant. It is hence meaningful to expand the terms above in powers of $\hat{\sigma}$. Going up to order $\hat{\sigma}^2$ one obtains for X' and X''

$$\begin{aligned} X'(u, u') &= D_A(u) \hat{\sigma}(u, u') D_B(u') \\ &\quad - \left[\int D_A(u) \hat{\sigma}(u, u') C_B(u', u'_1) \hat{\sigma}(u'_1, u_1) D_A(u_1) du_1 du'_1 + A \leftrightarrow B \right] \\ &\quad - \int C_A(u_1, u) \hat{\sigma}(u, u') C_B(u', u'_1) \hat{\sigma}(u'_1, u_1) du_1 du'_1 \end{aligned} \quad (28)$$

$$\begin{aligned} X''(u_1, u'_1, u_2, u'_2) &= - \left[D_A(u_1) \hat{\sigma}(u_1, u'_1) C_B(u'_1, u'_2) \hat{\sigma}(u_2, u'_2) D_A(u_2) + A \leftrightarrow B \right] \\ &\quad - C_A(u_1, u_2) \hat{\sigma}(u_1, u'_1) C_B(u'_1, u'_2) \hat{\sigma}(u_2, u'_2) \end{aligned} \quad (29)$$

At order $\hat{\sigma}^2$, the single scattering "exclusive" cross section $\tilde{\sigma}_1$ is given by

$$\begin{aligned} \tilde{\sigma}_1(u, u') &= D_A(u) \hat{\sigma}(u, u') D_B(u') \left[1 - \int D_A(u_1) \hat{\sigma}(u_1, u'_1) D_B(u'_1) du_1 du'_1 \right] \\ &\quad - \left[\int D_A(u) \hat{\sigma}(u, u') C_B(u', u'_1) \hat{\sigma}(u'_1, u_1) D_A(u_1) du_1 du'_1 + A \leftrightarrow B \right] \\ &\quad - \int C_A(u_1, u) \hat{\sigma}(u, u') C_B(u', u'_1) \hat{\sigma}(u'_1, u_1) du_1 du'_1 \end{aligned} \quad (30)$$

while the expression of the double scattering "exclusive" cross section $\tilde{\sigma}_2$ is

$$\begin{aligned} \tilde{\sigma}_2(u_1, u'_1, u_2, u'_2) &= \frac{1}{2!} \left\{ D_A(u_1) \hat{\sigma}(u_1, u'_1) D_B(u'_1) \cdot D_A(u_2) \hat{\sigma}(u_2, u'_2) D_B(u'_2) \right. \\ &\quad + \left[D_A(u) \hat{\sigma}(u_1, u'_1) C_B(u'_1, u'_2) \hat{\sigma}(u_2, u'_2) D_A(u_2) + A \leftrightarrow B \right] \\ &\quad \left. + C_A(u_1, u_2) \hat{\sigma}(u_1, u'_1) C_B(u'_1, u'_2) \hat{\sigma}(u_2, u'_2) \right\} \end{aligned} \quad (31)$$

One may immediately verify that

$$\begin{aligned} \int \tilde{\sigma}_1(u, u') du du' + 2! \int \tilde{\sigma}_2(u_1, u'_1, u_2, u'_2) du_1 du'_1 du_2 du'_2 &= \int D_A(u) \hat{\sigma}(u, u') D_B(u') du du' \\ &= \langle N \rangle \sigma_{hard} \equiv \sigma_S \end{aligned} \quad (32)$$

By restricting the phase space interval of the observed final state, $\tilde{\sigma}_1$ is well expressed by the term linear in $\hat{\sigma}$, which means that in that limit the probability of interaction is well approximated by the average of the distribution, while $\tilde{\sigma}_2$ is negligibly small. In other words the single parton scattering "exclusive" cross section is well represented by the single scattering expression of the simple QCD parton model. When the phase space volume is increased, the single parton scattering "exclusive" cross section becomes increasingly different from the prediction of the QCD parton model and, as it will be shown in the explicit examples of the next two paragraphs, the difference allows a direct measure of the importance of correlations. Notice that, by summing the expressions of $\tilde{\sigma}_1$ and $\tilde{\sigma}_2$ at order $\hat{\sigma}^2$ with the proper multiplicity factors one obtains, as shown above, the *inclusive* single parton scattering cross section, which is correctly given by the QCD parton model expression. By comparing the measured inclusive cross section with a sum of the measured "exclusive" cross sections, taken with the proper multiplicity factors, up to a given order in the number of collisions, one hence has a direct indication of the importance of higher order unitarity corrections in a given phase space interval.

To show how these ideas may be explicitly implemented we work out two explicit examples in the following paragraphs 4.1 and 4.2. The transverse parton coordinates are not accessible experimentally. On the other hand while, at small x , partons may not be correlated in momentum fraction, as conservation constraints may be diluted when the parton population is large, they are surely correlated in their transverse coordinates, since they are all confined inside the same hadron. A simplest possibility is hence to assume that the only relevant correlations are those in the transverse coordinates. To have some indication we will consider two different analytically solvable models of the parton densities in transverse space, the Gaussian and the exponential model, while neglecting in our two examples all effects of longitudinal correlations.

4.1. Gaussian model

One may assume Gaussian distributions for the parton densities and for the correlations:

$$\begin{aligned} D(x, b) &= G(x) f(b) \\ f(b) &= g(b, R^2) \\ C(x_1, x_2; b_1, b_2) &= G(x_1) G(x_2) h(b_1, b_2) \\ h(b_1, b_2) &= c \cdot g(B, R^2/2) \bar{h}(b, \lambda) \\ \bar{h}(b, \lambda) &= \eta \frac{d}{d\gamma} g(b, \lambda^2/\gamma) \Big|_{\gamma=1} \\ g(b, R^2) &= \frac{1}{\pi R^2} \exp(-b^2/R^2) \end{aligned} \quad (33)$$

where

$$\begin{aligned}
\mathbf{B} &= [\mathbf{b}_1 + \mathbf{b}_2]/2 \\
\mathbf{b} &= [\mathbf{b}_1 - \mathbf{b}_2]
\end{aligned}
\tag{34}$$

in such a way that the following relations hold

$$\begin{aligned}
\int d^2b g(b, R^2) &= 1, & \int d^2b_2 g(\mathbf{b}_1 - \mathbf{b}_2, R_1^2) g(\mathbf{b}_2, R_2^2) &= g(\mathbf{b}_1, R_1^2 + R_2^2) \\
\int d^2b h(b_1, b_2) d^2b &= 0
\end{aligned}
\tag{35}$$

The connection with the root mean square hadron radius $\langle r^2 \rangle \equiv \bar{R}^2$ is $R^2 = 2\bar{R}^2/3$. We define the correlation r_c as the value of b where the function $\bar{h}(b, \lambda)$ changes sign. In this case one hence has $r_c = \lambda$. In order to define unambiguously the correlation strength we choose the normalization

$$\int_{|b| \leq r_c} \bar{h}(b, \lambda^2) d^2b = 1,$$

since the integral over the whole b -space gives zero. With this choice $\eta = e$, where e is the Euler number. The integrations on the transverse variables of the various terms in Eq.s 30 and 31 give

$$\begin{aligned}
D_A \hat{\sigma} D_B &\rightarrow \int d^2b d^2\beta g_A(\mathbf{b} - \boldsymbol{\beta}, R_1^2) g_B(\mathbf{b}, R_2^2) = 1 \\
D_A \hat{\sigma} D_B \cdot D_A \hat{\sigma} D_B &\rightarrow \int d^2b_1 d^2b_2 d^2\beta g_A(\mathbf{b}_1 - \boldsymbol{\beta}, R_1^2) g_B(\mathbf{b}_1, R_2^2) g_A(\mathbf{b}_2 - \boldsymbol{\beta}, R_1^2) g_B(\mathbf{b}_2, R_2^2) \\
&= \frac{1}{2\pi(R_1^2 + R_2^2)} \\
D_A \hat{\sigma} C_B \hat{\sigma} D_A &\rightarrow \int d^2b_1 d^2b_2 d^2\beta g_A(\mathbf{b}_1 - \boldsymbol{\beta}, R_1^2) h_B(\mathbf{b}_1, \mathbf{b}_2) g_A(\mathbf{b}_2 - \boldsymbol{\beta}, R_1^2) \\
&= \frac{c e}{\pi} \frac{\lambda^2}{(2R_1^2 + \lambda^2)^2} \\
C_A \hat{\sigma} C_B \hat{\sigma} &\rightarrow \int d^2b_1 d^2b_2 d^2\beta h_A(\mathbf{b}_1 - \boldsymbol{\beta}, \mathbf{b}_2 - \boldsymbol{\beta}) h_B(\mathbf{b}_1, \mathbf{b}_2) \\
&= \frac{c^2}{4\pi} \frac{e^2}{\lambda^2}
\end{aligned}
\tag{36}$$

the "exclusive" cross sections $\tilde{\sigma}_1$ and of $\tilde{\sigma}_2$ are hence explicitly expressed in this case in terms of the QCD-parton model single scattering cross section σ_S , the transverse hadron size R and the correlations parameters c and λ :

$$\begin{aligned}
\frac{d\tilde{\sigma}_1}{dyd\mathbf{p}_t} &= \frac{d\sigma_S}{dyd\mathbf{p}_t} \times \left[1 - \frac{\sigma_S}{4\pi R^2} - ce \frac{2\sigma_S \lambda^2}{\pi(2R^2 + \lambda^2)^2} - c^2 e^2 \frac{\sigma_S}{4\pi \lambda^2} \right] \\
&= \frac{d\sigma_S}{dyd\mathbf{p}_t} \times \left[1 - \frac{\sigma_S}{\sigma_{eff}} \right] \\
\frac{d\tilde{\sigma}_2}{dy_1 dy_2 d\mathbf{p}_{t1} d\mathbf{p}_{t2}} &= \frac{1}{2} \frac{d\sigma_S}{dy_1 d\mathbf{p}_{t1}} \frac{d\sigma_S}{dy_2 d\mathbf{p}_{t2}} \times \left[\frac{1}{4\pi R^2} + ce \frac{2\lambda^2}{\pi(2R^2 + \lambda^2)^2} + c^2 e^2 \frac{1}{4\pi \lambda^2} \right] \\
&= \frac{1}{2\sigma_{eff}} \frac{d\sigma_S}{dy_1 d\mathbf{p}_{t1}} \frac{d\sigma_S}{dy_2 d\mathbf{p}_{t2}} \equiv \frac{d\sigma_D}{dy_1 dy_2 d\mathbf{p}_{t1} d\mathbf{p}_{t2}}
\end{aligned} \tag{37}$$

where σ_{eff} is the scale factor which characterizes the inclusive rate of double parton collisions. As a function of the root mean hadron radius $\bar{R}$, of the correlation length r_c and of the correlation strength c , the expression is

$$\frac{1}{\sigma_{eff}} = \frac{3}{8\pi \bar{R}^2} \left\{ 1 + ce \cdot 16 \frac{3s^2}{(4 + 3s^2)^2} + c^2 e^2 \cdot \frac{2}{3s^2} \right\} \tag{38}$$

where $s \equiv r_c/\bar{R}$. Notice that neglecting correlation in fractional momenta, in each phase space interval $\tilde{\sigma}_1$ is different from σ_S only by the overall normalization (related to σ_{eff} as shown above). At this order in $\tilde{\sigma}$, $\tilde{\sigma}_2$ coincides with the expression of the inclusive double parton scattering cross section while, in this case, the scale factor σ_{eff} does not depend on x .

4.2. Exponential model

A different option is to assume an exponential for the three dimensional parton density

$$\begin{aligned}
\rho(r) &= \frac{\mu^3}{8\pi} e^{-\mu r}, \quad \int \rho(r) d^3r = 1 \\
D(x, b) &= G(x) f(b) \\
f(b) &= \int \rho(r) dz = \frac{1}{4\pi} \mu^3 b K_1(\mu b)
\end{aligned} \tag{39}$$

where $\mathbf{b}$ is the transverse component of $\mathbf{r}$ and K_1 is the modified Bessel function of the second kind. It may be seen that the two distributions Gaussian and exponential are formally connected. If one performs a linear combination of the Gaussian distributions (33) with appropriate weight one ends up with the expression (39):

$$\int_0^\infty \frac{\mu^4}{2(2\lambda)^3} e^{-\mu^2/4\lambda} g(b, 1/\lambda) d\lambda = \frac{1}{4\pi} \mu^3 b K_1(\mu b) \tag{40}$$

For the next calculations it is convenient to take the Fourier transform of the transverse parton density

$$\begin{aligned}
\rho(b) &= \frac{1}{(2\pi)^2} \int e^{-i\mathbf{p}_t \cdot \mathbf{b}} \tilde{\rho}(p_t) d^2 p_t \\
\tilde{\rho}(p_t) &= \frac{\mu^4}{(\mu^2 + p_t^2)^2}
\end{aligned} \tag{41}$$

One may assume for the correlation the expression

$$\begin{aligned}
C(x_1, x_2; b_1, b_2) &= G(x_1)G(x_2)h(b_1, b_2) \\
h(b_1, b_2) &= c \cdot f(B)\bar{h}(b) \\
\bar{h}(b) &= \frac{1}{(2\pi)^2} \int e^{-i\mathbf{p}_t \cdot \mathbf{b}} \tilde{h}(p_t) d^2 p_t \\
\tilde{h}(p_t) &= \eta \frac{d}{d\gamma} \frac{\kappa^4 \gamma^2}{(\kappa^2 + p_t^2)^2} \Big|_{\gamma=1} = 2\eta \frac{\kappa^4 p_t^2}{(p_t^2 + \kappa^2)^3}
\end{aligned} \tag{42}$$

in such a way that

$$\tilde{h}(0) = \int \bar{h}(b) d^2 b = 0 \tag{43}$$

As a function of the coordinates the expression is

$$\bar{h}(b) = \frac{\eta}{2\pi} \left\{ \kappa^3 b K_1(\kappa b) - \frac{1}{4} \kappa^4 b^2 K_2(\kappa b) \right\} \tag{44}$$

where K_1 and K_2 are modified Bessel functions. The root mean square value correlation term is performed in the same way as in of the density is $\langle r^2 \rangle \equiv \bar{R}^2 = 12/\mu^2$ and, like in the previous case, defining the correlation length as the value of b where the function $\bar{h}(b)$ changes sign, one has $r_c = x_0/\kappa$, where $x_0 \simeq 2.386$ solves the equation

$$4x_0 K_1(x_0) = x_0^2 K_2(x_0) \tag{45}$$

The normalization of the correlation function is performed in the same way as in the Gaussian model, it gives now $\eta = 3.456$. The integrations on the transverse variables of the various terms in Eq.s 30 and 31 give

$$\begin{aligned}
D_A \hat{\sigma} D_B &\rightarrow \int d^2 b d^2 \beta f_A(\mathbf{b} - \beta) f_B(\mathbf{b}) = 1 \\
D_A \hat{\sigma} D_B \cdot D_A \hat{\sigma} D_B &\rightarrow \int d^2 b_1 d^2 b_2 d^2 \beta f_A(\mathbf{b}_1 - \beta) f_B(\mathbf{b}_1) f_A(\mathbf{b}_2 - \beta) f_B(\mathbf{b}_2) \\
&= \frac{3}{7\pi \bar{R}^2} \\
D_A \hat{\sigma} C_B \hat{\sigma} D_A &\rightarrow \int d^2 b_1 d^2 b_2 d^2 \beta f_A(\mathbf{b}_1 - \beta) \bar{h}_B(\mathbf{b}_1, \mathbf{b}_2) f_A(\mathbf{b}_2 - \beta) \\
&= c \frac{3\eta}{\pi \bar{R}^2} F\left[\frac{x_0^2 \bar{R}^2}{12r_c^2}\right] \\
F(a) &\equiv \frac{3 + 44a - 36a^2 - 12a^3 + a^4 + 12a(2 + 3a)\ln(a)}{3(a - 1)^6 a} \\
C_A \hat{\sigma} C_B \hat{\sigma} &\rightarrow \int d^2 b_1 d^2 b_2 d^2 \beta \bar{h}_A(\mathbf{b}_1 - \beta, \mathbf{b}_2 - \beta) \bar{h}_B(\mathbf{b}_1, \mathbf{b}_2) \\
&= c^2 \eta^2 \frac{1}{30\pi} \frac{x_0^2}{r_c^2}
\end{aligned} \tag{46}$$

Notice that the function $F(a)$ is regular near $a = 1$, the expansion being

$$F(a) = \frac{1}{15} - \frac{a - 1}{7} + 3 \frac{(a - 1)^2}{14} - 5 \frac{(a - 1)^3}{18} + \mathcal{O}((a - 1)^4) \tag{47}$$

As a function of the root mean square hadron radius $\bar{R}$, of the ratio $s \equiv r_c/\bar{R}$ and of the correlation strength c , the expression of the effective cross section is

$$\frac{1}{\sigma_{eff}} = \frac{3}{7\pi \bar{R}^2} \left\{ 1 + c\eta \cdot 14 \cdot F\left(\frac{1}{12} \frac{x_0^2}{s^2}\right) + c^2 \eta^2 \cdot \frac{7}{30} \left(\frac{1}{3} \frac{x_0^2}{s^2}\right) \right\} \tag{48}$$

One may hence obtain information on the correlations by comparing the measured "exclusive" single scattering cross section with the measured inclusive single scattering cross section:

$$\frac{d\tilde{\sigma}_1}{dy d\mathbf{p}_t} - \frac{d\sigma_S}{dy d\mathbf{p}_t} = \frac{d\sigma_S}{dy d\mathbf{p}_t} \frac{\sigma_S}{\sigma_{eff}} \tag{49}$$

The cases considered here are relevant for a phase space window where multiple collisions, of order higher than two, do not give important contributions; condition which may be always achieved by taking, for example, the lower cutoff in p_t large enough. Notice that the importance of higher order scattering terms, namely of unitarity corrections, is controlled irrespectively of any simplifying hypothesis on correlations, by verifying the validity of the sum rule Eq.32.

The relation above allows to obtain the value of σ_{eff} . By subtracting from the effective cross section the term $3/(7\pi \bar{R}^2)$, or $3/(8\pi \bar{R}^2)$, depending on the prejudice on the parton density in transverse space, one obtains a direct information on correlations. Notice that, although the two options considered, the Gaussian and the exponential, correspond to rather different parton densities, the quantity to be subtracted is basically the same in the two cases and it is essentially

determined only by the value of the mean square hadron radius $\bar{R}^2$. Indications on the value of $\bar{R}^2$ are obtained by the measurement of the generalized parton distributions[22][23]. In the case of gluons one may estimate $\bar{R}^2 \simeq .42 \text{ fm}^2$ [24].

One should point out that the validity of Eq.49 is easily tested, since the dependence of the difference in the right hand side of the equation, integrated in a given phase space interval, on the rapidity window and on the lower cutoff in p_t has to be the same as the dependence of σ_S^2 , integrated in the same phase space interval. So, if the sum rule of Eq.32 is satisfied while the relation in Eq.49 is not, one may safely conclude that one has direct evidence of longitudinal correlations in the hadron structure.

The analysis may be repeated by including higher order multiple scattering terms in Eq.30 and in the following. In such a case, one may write relations similar to Eq.49, linking higher order multiple scattering inclusive cross sections with various combinations of "exclusive" cross sections. Independent constraints on the correlation parameters may hence be established, hopefully allowing an unambiguous identification of the correlation parameters.

5. CONCLUDING REMARKS

MPI are originated by the large flux of partons in high energy hadronic collisions. The dominant contribution to the process is hence due to the terms which maximize the incoming parton flux. Given the number of partonic collisions, the incoming flux is maximized by the set of terms where elementary hard interactions are all disconnected one from another or, more precisely, where the connection is only due to soft exchanges. Being the interactions uncorrelated with respect to hard dynamics, all correlations observable in the final state are originated by the initial state flux, namely by the hadron structure. In such a scheme all partonic collisions add incoherently in the final cross section, leading to a purely probabilistic picture of the process. It is hence natural to identify two different classes of hard cross sections, the inclusive cross section, related to the moments of the distribution, and the "exclusive" cross sections, related to the different terms of the probability distribution in the number of hard collisions.

A relevant property is that the picture of the interaction is probabilistic in each phase space interval, in such a way that one may associate a different probability distributions to each different phase space choice to observe the final state. Including the possibility to select events which are fully or only partially contained in the given phase space interval. Any choice identifies an interval in p_t and restricts in a precise way the limits in x of the initial state, which identifies the domain of definition of the probability distribution of multiple collisions. One should point out that the same restricted intervals define also the integration range of the integrated terms, which appear in the "exclusive" differential cross sections (e.g. the integrations in Eq.30). Integrated terms appear in the "exclusive" differential cross sections because of the normalization of the probability distribution, their origin may be traced back to the virtual corrections of the process. Normalization hence fixes unambiguously the integration limits of the virtual terms, which must coincide with the kinematical limits imposed to the real terms by the choice adopted to select the final state.

Another relevant point is that, as proved in the appendix, the components of the hadron structure, namely the terms D and C_n , are well defined quantities in the picture of the interaction,

as they do not mix when the selection choice of the final state is modified. The effect, to modify the selection choice of the final state, is in fact only to change the integration domain of each term, namely to probe the multi-parton structure of the hadron in different domains in x and Q^2 .

As it was pointed out, the inclusive cross sections can provide a direct information on the correlations of the multi-parton structure of the hadron, while to evaluate the "exclusive" cross sections one needs, in principle, an infinite non perturbative information. The expansion of the "exclusive" cross sections in the number of parton collisions allows however to identify the dominant terms and to isolate the leading contributions due to the presence of correlations. Explicit relations between inclusive and exclusive cross sections may hence be written at each order of the expansion in the number of collisions, while the importance of higher order terms may be controlled by selecting the phase space interval where to observe the final state. The comparison between experimentally measured inclusive and "exclusive" cross sections, at a given order in the number of collisions, provides a model independent tool to judge on the importance of higher order unitarity corrections (namely by checking the validity of sum rules like Eq.32).

In the present paper we have worked out in details, at the lowest non trivial order, the simplest case where only the correlations in the transverse coordinates are taken into account. Of course correlations will depend also on fractional momenta and on the different kinds of partons considered (gluons and quarks, distinguishing different flavors and the valence from the sea). To some extent the different kinds of partons involved may be selected by looking at definite final states produced and at the corresponding values of momentum fraction involved (a prompt photon plus five jets at low x are likely to be generated by a triple scattering involving a quark and five gluons etc.). The information on correlations will hence to be related to the different kinds of initial state partons involved in the interaction.

While the importance of higher order unitarity corrections is tested, in a given phase space interval, by verifying the sum rule in Eq.32 (and similar when including a higher number of collisions) terms with correlations are isolated by using relations like Eq.49, which allow to obtain the value of the effective cross section and of similar scale factors for higher orders unitarity corrections. The importance of correlations is tested by comparing with the expectation in the uncorrelated case and, in particular, longitudinal momentum correlations will show up with the dependence of the effective cross section (and of the similar scale factors for higher orders unitarity corrections) on the rapidity window. With respect to longitudinal correlations one should anyhow point out that, while indications on the effects of longitudinal correlations may be obtained along these lines in pp collisions, a model independent separation of longitudinal and transverse correlations may be obtained by looking at MPI in the case of hadron-nucleus collisions[21].

Acknowledgments

One of us (D.T.) thanks Mark Strikman and Ted Rogers for useful discussions.

APPENDIX

As already mentioned in the main text the "exclusive" cross sections are defined with respect to intervals of variation of the kinematical variables, which are to some extent arbitrary. Even

though the result may look intuitive, we like to analyze formally what happens when varying the intervals of the kinematical variables: Starting with the formalism stated in eq. (8) we call $\mathcal{U}$ the field of variation of u and divide it into two mutually exclusive parts $\mathcal{U} = \mathcal{R} \cup \mathcal{S}$. Correspondingly it is convenient to denote by r the variable u when it varies within $\mathcal{R}$ and s when it varies within $\mathcal{S}$. Then for the exclusive distribution within $\mathcal{R}$ we have:

$$\tilde{W}_1(r_1) = W_1(r_1) + \int_{\mathcal{S}} W_2(r_1, s) ds + \frac{1}{2} \int_{\mathcal{S}} W_3(r_1, s, s') ds ds' + \frac{1}{3!} \int_{\mathcal{S}} W_4(r_1, s, s', s'') ds ds' ds'' + \dots$$

$$\tilde{W}_2(r_1, r_2) = W_2(r_1, r_2) + \int_{\mathcal{S}} W_3(r_1, r_2, s) ds + \frac{1}{2} \int_{\mathcal{S}} W_4(r_1, r_2, s, s') ds ds' + \dots$$

These relations may be summarized by means of the following procedure: Let us define the projectors $R(u) = 1, S(u) = 0$ for $u \in \mathcal{R}$ $R(u) = 0, S(u) = 1$ for $u \in \mathcal{S}$, $R + S = 1$; and the generating functional $\tilde{\mathcal{Z}}[J] = \mathcal{Z}[R \cdot J + S]$. Then expanding $\tilde{\mathcal{Z}}[J]$ as in eq (8) and using the symmetry of the W_n it is easy to attain the result:

$$\tilde{\mathcal{Z}}[J] = \sum_n \int_{\mathcal{R}} J(r_1) \dots J(r_n) \tilde{W}_n(r_1 \dots r_n) dr_1 \dots dr_n$$

Now we perform the same substitution in the logarithm of the generating functional $\tilde{\mathcal{F}}[J] = \mathcal{F}[R \cdot J + S]$ and using $1 - S = R$ the projector R is found in front of every integrand, so the overall effect is a restriction in the integration domain and the equivalent of eq. (10) is easily found:

$$\tilde{\mathcal{F}}[J] = \int_{\mathcal{R}} D_1(r) [J(r) - 1] du + \sum_{n=2}^{\infty} \frac{1}{n!} \int_{\mathcal{R}} C_n(r_1 \dots r_n) [J(r_1) - 1] \dots [J(r_n) - 1] dr_1 \dots dr_n$$

The conclusion is hence that the inclusive quantities are not modified by the variation of the interval where the kinematical variables are defined, on the contrary the variation of the exclusive quantities can be worked out. With respect to the previous formula the difference arises only from the different integration domain of the terms D and C_n .

-
- [1] T. Sjostrand and M. van Zijl, Phys. Rev. D **36**, 2019 (1987).
 - [2] D. E. Acosta *et al.* [CDF Collaboration], Phys. Rev. D **70**, 072002 (2004) [arXiv:hep-ex/0404004].
 - [3] T. Sjostrand, S. Mrenna and P. Skands, JHEP **0605**, 026 (2006) [arXiv:hep-ph/0603175].
 - [4] M. Bahr, S. Gieseke and M. H. Seymour, JHEP **0807**, 076 (2008) [arXiv:0803.3633 [hep-ph]].
 - [5] T. Akesson *et al.* [Axial Field Spectrometer Collaboration], Z. Phys. C **34**, 163 (1987).
 - [6] J. Alitti *et al.* [UA2 Collaboration], Phys. Lett. B **268**, 145 (1991).
 - [7] F. Abe *et al.* [CDF Collaboration], Phys. Rev. Lett. **79**, 584 (1997).
 - [8] F. Abe *et al.* [CDF Collaboration], Phys. Rev. D **56**, 3811 (1997).
 - [9] G. Calucci and D. Treleani, Phys. Rev. D **57**, 503 (1998) [arXiv:hep-ph/9707389].
 - [10] A. Capella, J. Tran Thanh Van and J. Kwiecinski, Phys. Rev. Lett. **58**, 2015 (1987).
 - [11] L. Ametller and D. Treleani, Int. J. Mod. Phys. A **3**, 521 (1988).
 - [12] T. C. Rogers, A. M. Stasto and M. I. Strikman, Phys. Rev. D **77**, 114009 (2008) [arXiv:0801.0303 [hep-ph]].

- [13] J. M. Butterworth, J. R. Forshaw and M. H. Seymour, *Z. Phys. C* **72**, 637 (1996) [arXiv:hep-ph/9601371].
- [14] T. Sjostrand, arXiv:0809.0303 [hep-ph].
- [15] B. Blaettel, G. Baym, L. L. Frankfurt, H. Heiselberg and M. Strikman, *Phys. Rev. D* **47**, 2761 (1993).
- [16] E. Gotsman, E. Levin and U. Maor, arXiv:0708.1506 [hep-ph].
- [17] L. Frankfurt, M. Strikman, D. Treleani and C. Weiss, arXiv:0808.0182 [hep-ph].
- [18] D. Treleani, arXiv:0808.2656 [hep-ph].
- [19] G. Calucci and D. Treleani, *Int. J. Mod. Phys. A* **6**, 4375 (1991).
- [20] A. Białas, M. Bleszynski and W. Czyz, *Nucl. Phys. B* **111**, 461 (1976).
- [21] M. Strikman and D. Treleani, *Phys. Rev. Lett.* **88**, 031801 (2002) [arXiv:hep-ph/0111468].
- [22] D. Mueller, D. Robaschik, B. Geyer, F. M. Dittes and J. Horejsi, *Fortsch. Phys.* **42**, 101 (1994) [arXiv:hep-ph/9812448].
- [23] A. V. Belitsky and A. V. Radyushkin, *Phys. Rept.* **418**, 1 (2005) [arXiv:hep-ph/0504030].
- [24] L. Frankfurt, M. Strikman and C. Weiss, *Phys. Rev. D* **69**, 114010 (2004) [arXiv:hep-ph/0311231].